

Order of superconductive phase transition

2 H.Kleinert

Institut für Theoretische Physik,
Freie Universität Berlin,
Arnimallee 14, D–14195 Berlin, Germany

Received December 26, 2004

3 On the occasion of Reinhard Folk's 60th birthday, I give a brief review of
4 the theoretical progress in understanding the critical properties of super-
5 conductors. I point out the theoretical difficulties in finding a second-order
6 transition in the Ginzburg-Landau Model with $O(N)$ -symmetry in $4 - \epsilon$ Di-
7 mensions, and the success in predicting the existence and location of a
8 tricritical point with the help of a dual disorder theory.

9 **Key words:**

10 **PACS:**

11 1. Introduction

12 Among the many important contributions made by Reinhard Folk in the field
13 of critical phenomena is his work on the properties of the superconducting phase
14 transitions [1,2], where he discussed the effect of two-loop contributions on the renor-
15 malization flow. To point out its significance, let me briefly recall the historic back-
16 ground of the problem studied by him with various collaborators, most prominently
17 with Yuriy Holovatch.

18 Until thirty years ago, the superconductive phase transition was always assumed
19 to be of second order. All critical exponents had the mean-field values required by
20 the Ginzburg-Landau model. In 1974, however, it was noted by Halperin, Lubensky,
21 and Ma [3] (see also [4]) that the renormalization group treatment of the Ginzburg-
22 Landau model of superconductivity did not produce a critical point. A similar ob-
23 servation had been published earlier in four-dimensional quantum field theory by
24 Coleman and Weinberg [5] (working at the same institute). This was interpreted
25 as an indication that thermal fluctuations which, in general, cause only small devi-
26 ations of the critical exponents from their mean-field values, would here have the
27 more dramatic effect of driving the superconductive transition first order. At that
28 time, the theoretical result could not be tested by experiment. The reason is that in

old-fashioned superconductors, the temperature interval around the critical point, in which fluctuations become important, the so-called *Ginzburg interval* [6], is extremely narrow, lying in the nK-regime. This explained the observed mean-field behavior. The critical regime had always been too small to be resolved. See the textbooks [7,8] for details.

To probe the Ginzburg interval and confirm the second-order nature of the transition, one had to rely on Monte-Carlo simulations [9] and a theoretical analogy with smectic-nematic transitions in liquid crystals [10], which is also described by a Ginzburg-Landau model, albeit with a different interpretation of the fields [11] (the so-called Landau-De Gennes model).

The situation has changed since the discovery of high- T_c superconductors, where the Ginzburg interval is large, up to several K. Recent experiments [12] clearly show a critical behavior associated with a charged critical point, in which both the coherence length ξ and the magnetic penetration depth λ grow with the same critical exponent $\nu_\lambda = \nu_\xi \approx 2/3$ [13].

The confusing critical properties of the Ginzburg-Landau model certainly required further investigation, in particular it appeared necessary to calculate higher loop approximations. To appreciate the problem, let us first recall what makes the first-order result of [3] questionable: The simplest argument leading to the first order is based on a fluctuation-corrected mean-field theory obtained in three dimensions by integrating out exactly the gauge field at fixed $|\phi|$. The resulting trace-log can be evaluated exactly, and produces a cubic term $\propto -|\phi|^3$, which makes the transition first-order. The trouble with this argument is that in the type II regime, fluctuations will produce vortex lines. At the core of each vortex line, the order field ϕ vanishes, which invalidates the assumption of a fixed $|\phi|$ in the above calculation.

The first direct theoretical evidence for a second-order phase transition did not come from the Ginzburg-Landau model but from a dual formulation of it on a lattice which I set up about 20 years ago [14]. Its properties are discussed in great detail in the textbook [7]. The dual formulation contains no massless gauge field, and this avoids the above problem coming from the zeroth in the order field ϕ . My dual model predicted definitely a second-order transition in the type-II regime characterized by a Ginzburg parameter $\kappa > 1/\sqrt{2}$. This parameter measures the ratio between magnetic penetration depth and coherence length in the superconductor.

The dual theory contains a different field ψ which is a *disorder field*. The associated Feynman diagrams are direct pictures of the vortex loops, whose density is $|\psi|^2$. This field has a quartic self-interaction accounting for their short-range repulsion, and a gauge-coupling to a *massive* vector field \vec{h} representing the fluctuating magnetic induction in the superconducting phase. Due to the mass term, the field \vec{h} can be integrated out and the assumption of a constant mean field $|\psi|$ presents no problem, leading to a Landau-like expansion of the free energy containing terms $|\psi|^4$, $|\psi|^6$, etc. The $|\psi|^4$ term turned out to be proportional to $\kappa - \kappa_t$, with [14]

$$\kappa_t \approx \frac{3\sqrt{3}}{2\pi} \sqrt{1 - \frac{4}{9} \left(\frac{\pi}{3}\right)^4} \equiv \frac{0.798}{\sqrt{2}}. \quad (1)$$

1 For $\kappa < \kappa_t$, vortices attract each other on the average, and the transition is of first
2 order, whereas for $\kappa > \kappa_t$, they repel each other and the transition is of second order.
3 Thus, the mean-field approximation in the disorder theory suggests a second-order
4 transition in the type II regime of repulsive vortex lines and a first-order transition
5 in the type I regime of attractive vortex lines.

6 The point $\kappa = \kappa_t$ is a *tricritical point*. Its existence has been confirmed by Monte
7 Carlo simulations on a lattice in [15], but initially at quite a different value $\kappa_t \approx$
8 $0.38/\sqrt{2}$. Only recently, with the availability of much better simulation techniques,
9 has my 1982 prediction been confirmed with amazing accuracy [16].

10 In order to satisfactorily understand the transition, one should be able to ex-
11 plain the second order of the transition in the type-II regime without the duality
12 argument. Thus we must find the effective potential of the Ginzburg-Landau model
13 as accurately as possible. So far, only the two-loop effective potential is known for
14 $N/2$ complex fields in $4 - \epsilon$ dimensions, which I calculated together with my collab-
15 orator Van den Bossche [17]. The result is given in sections 3 and 4. The case $N = 2$
16 concerns the physical situation of the Ginzburg-Landau model of superconductivity.

17 The involved calculations became possible due to recent progress made in evalu-
18 ation techniques of Feynman diagrams in $4 - \epsilon$ dimensions. In particular, two-loop
19 Feynman diagrams with unequal masses of the internal lines have become available
20 analytically [18,19]. Moreover, the full ϵ -expansion of the so-called sunset diagram is
21 now known [20]. This is important for critical phenomena in $4 - \epsilon$ dimensions since,
22 for instance, a three-loop calculation requires the knowledge of all terms of order ϵ
23 of the two-loop diagrams.

24 Unfortunately, the renormalization flow of the coupling constants in $4 - \epsilon$ dimen-
25 sions does not improve much over the one-loop calculation of [3] when going to two
26 loops [21,22]. Apparently, a direct three-dimensional approach is more promising to
27 lead to a second-order transition for $N = 2$ [23,24]. The absence of a charged fixed
28 point for $N = 2$ in $4 - \epsilon$ dimensions seems thus to be a specific weakness of the
29 ϵ -expansion, although important progress has been made in [1]: There, a [1/1] Padé-
30 Borel resummation of the two beta functions (associated with the electric charge and
31 with the self-coupling of the scalar field) has indeed lead to a desired IR stable fixed
32 point for $N = 2$. The resummation at such a low order is, however, not very reliable
33 and the result should be considered as fortuitous.

34 After all this work it is not even certain that a higher-loop renormalization group
35 analysis can really lead to a critical point. In my opinion, the perturbation expansion
36 to low orders knows too little about the vortex fluctuations, which should be included
37 explicitly into the path integral [25], and which the dual theory takes optimally into
38 account. In ϕ^4 -theory without gauge field, this seems to be unnecessary for reasons
39 which are not yet understood. The omission is only justified there by the success in
40 predicting the experimental critical exponents in these systems [8].

41 We hope that eventually *variational perturbation theory* will be capable of locati-
42 ng a fixed point of the Ginzburg-Landau model, thus allowing us to extract physical
43 values independently of [1]. This theory has been developed and discussed in detail
44 in [8,26–30] and it has proven to be a powerful tool for determining critical exponents

in three [28] as well as in $4 - \varepsilon$ dimensions [30] of pure ϕ^4 -theories. Variational perturbation theory is a procedure which allows us to determine resummed quantities from a strong-coupling limit of divergent expansions in powers of the bare coupling constant. It no more requires Padé or Padé-Borel resummation. In particular, there is no freedom in selecting different Padé approximants which yield different results at higher orders than [1/1]. Variational perturbation theory has yielded the most precise prediction [26] $\alpha \approx -0.0129 \pm 0.0006$ for the best-measured critical exponent α governing the singularity of the specific heat of superfluid helium. To avoid the smear-out of the critical regime by gravity over 10^{-6} K and to determine the critical temperature up to 2nK, the experiment has been performed in the satellite [31] and gave $\alpha = -0.0127 \pm 0.0003$.

Recently, we have applied variational perturbation theory to a new range of problems: the determination of amplitude ratios in three dimensions of the $O(N)$ -model [32]. We based these calculations on expansions of the critical exponents obtained by a certain regularization method [33], in which analytic regularization is applied with a formal minimal subtraction in $4 - \varepsilon$ dimensions, but inserting at the end $\varepsilon = 1$ without invoking further ε -expansions. Our results were consistent with Padé-Borel methods. Due to its power, we believe that variational perturbation theory will eventually succeed in producing the desired zeros in the beta functions to yield an IR-stable charged fixed point.

When calculating the effective potential to be given in sections 3 and 4, it was useful to confirm the agreement of our renormalization constants in [17] with those obtained by Reinhard Folk and his collaborator Kolnberger of 1992 [21]. A student of mine had done this calculation eight years earlier for his M.S. thesis [22], but one of his coefficients contained an error, which was corrected in Folk's work.

Let us also mention that the effective potential given in sections 3 and 4 may be seen as an extension to a two-coupling-constants problem studied by Brézin et al. [34] (see also [35]), who give the ε -expansion of the equation of state of the N -components ϕ^4 theory without a gauge field up to two loops [36].

2. Ginzburg-Landau model

The Lagrangian density contains $N/2$ complex scalar fields ϕ coupled to the abelian fields A_μ and reads, with a covariant gauge fixing,

$$\mathcal{L} = |D\phi|^2 + m^2\phi^2 + \frac{g}{3!}|\phi|^4 + \frac{1}{4}F_{\mu\nu}^2 + \frac{1}{2\alpha}(\partial_\mu A_\mu)^2, \quad (2)$$

where $D_\mu = \partial_\mu - ieA_\mu$ is the covariant derivative, $F_{\mu\nu}$ is the usual field-strength tensor, and α is a gauge parameter. In principle, there are also ghost fields which, however, decouple in the symmetric phase and remain massless. Working in dimensional regularization they do not contribute to the energy due to Veltman's rule $\int d^D p p^\alpha = 0$ for all α [8]. The effective potential will be obtained using the so-called *background-field method* of DeWitt [38]. We shift the scalar field by an unknown constant Φ : $\phi \rightarrow \Phi + \phi$. This generates new vertices. To simplify the calculation,

we shall use throughout the Landau gauge $\alpha \rightarrow 0$. This reduces the number of Feynman diagrams and, since $\alpha = 0$ enforces $\partial_\mu A_\mu \equiv 0$ at the Lagrangian level, removes a possible mixing of $A_\mu \phi$ and $A_\mu \phi^\dagger$ terms, thus decoupling scalar and gauge propagators. It is further advantageous to use real fields, defining

$$\phi = \frac{1}{\sqrt{2}}(\phi_1 + i\phi_2), \quad \Phi = \frac{1}{\sqrt{2}}(\Phi_1 + i\Phi_2). \quad (3)$$

Then the Lagrangian has the expansion around the background field:

$$\begin{aligned} \mathcal{L} = \mathcal{L}_0 + \frac{1}{2}\phi[G^T P^T + G^L P^L]\phi + \frac{1}{2}A_\mu D^T P_{\mu\nu}^T A_\nu + \frac{g}{4!}[(\phi^2)^2 + 4\Phi\phi(\phi^2)] \\ + e^2 A^2 \Phi \phi + \frac{1}{2}e^2 A_\mu^2 \phi^2 + eA_\mu(\phi_2 \partial_\mu \phi_1 - \phi_1 \partial_\mu \phi_2), \end{aligned} \quad (4)$$

where ϕ and Φ are now N components real fields written as two-dimensional iso-vectors $\phi = (\phi_1, \phi_2)$ and $\Phi = (\Phi_1, \Phi_2)$. The notation is:

$$\mathcal{L}_0 = \frac{1}{2}m^2\Phi^2 + \frac{g}{4!}\Phi^4, \quad (5)$$

$$G^T \equiv -\partial^2 + m_T^2 = -\partial^2 + m^2 + \frac{g}{3!}\Phi^2, \quad G^L \equiv -\partial^2 + m_L^2 = -\partial^2 + m^2 + \frac{g}{2}\Phi^2, \quad (6)$$

$$D^T \equiv -\partial^2 + m_\gamma^2 = -\partial^2 + e^2\Phi^2, \quad (7)$$

$$P_{ij}^T = \delta_{ij} - \frac{\Phi_i \Phi_j}{\Phi^2}, \quad P_{ij}^L = \frac{\Phi_i \Phi_j}{\Phi^2}, \quad P_{\mu\nu}^T = \delta_{\mu\nu} - \frac{\partial_\mu \partial_\nu}{\partial^2}, \quad (8)$$

- 1 with G^T, G^L being the transverse and longitudinal inverse propagators of the scalar
 2 field, and $D_{\mu\nu}^T$ is the inverse transverse propagator of the photon field. The transver-
 3 sality of the latter is due to the Landau gauge. Note that there is no term $eA_\mu \partial_\mu (\Phi_2 \phi_1 -$
 4 $\Phi_1 \phi_2)$ which would mix vector and scalar propagators.

So far, all quantities (fields, coupling constants, and masses) are bare quantities. Up to the second order in the loop expansion, divergences show up as poles in ε up to the order $1/\varepsilon^2$. They have to be removed to have a finite limit $\varepsilon \rightarrow 0$. This is achieved by the renormalization constants

$$\begin{aligned} \phi = \phi_r \sqrt{Z_\phi}, \quad A_\mu = A_{\mu r} \sqrt{Z_A}, \\ m^2 = m_r^2 \frac{Z_{m^2}}{Z_\phi}, \quad g = g_r \mu^\varepsilon \frac{Z_g}{Z_\phi^2}, \quad e = e_r \mu^{\varepsilon/2} \frac{Z_e}{Z_\phi \sqrt{Z_A}} = \frac{e_r}{\sqrt{Z_A}} \mu^{\varepsilon/2}, \end{aligned} \quad (9)$$

- 5 where in the last equation, the relation $Z_e = Z_\phi$ has been used, which is a con-
 6 sequence of a Ward identity. Intuitively, it comes from the requirement $D_\mu \phi \rightarrow$
 7 $\sqrt{Z_\phi} D_\mu \phi_r$ for the covariant derivative. In the above equations, the bare quantities are
 8 on the left-hand-side and the renormalized ones are on the right-hand-side, indicated
 9 by the subscript ‘‘r’’. Note that the vacuum energy requires a special treatment [37].

We now state the result for the effective potential derived in [17] up to two loops. It has the general form

$$V = V^{(0)} + \frac{\hbar}{(4\pi)^2} [V^{(1,0)} + \varepsilon V^{(1,\varepsilon)}] + \left[\frac{\hbar}{(4\pi)^2} \right]^2 V^{(2)}. \quad (10)$$

- 10 The expansion terms will be given in the following section.

3. Renormalized effective potential to zero- and one-loop order

The zero-loop effective potential is trivial:

$$V(l=0) = \frac{1}{2}m^2\Phi^2 + \frac{g}{4!}\mu^\varepsilon\Phi^4 + \frac{m^4}{g\mu^\varepsilon}. \quad (11)$$

The renormalized one-loop potential is simply a combination of the previous bare zero-order and the trace-log terms:

$$V(l=1) = \frac{1}{2}m^2\Phi^2 Z_{m^2}^{(1)} + \frac{g}{4!}\mu^\varepsilon\Phi^4 Z_g^{(1)} + \frac{m^4}{g\mu^\varepsilon} Z_v^{(1)} + \frac{\Gamma(1-D/2)}{(4\pi)^{D/2}} \frac{1}{D} \mu^{-\varepsilon} \\ \times \left[(N-1)(m_T^2)^2 \left(\frac{\mu^2}{m_T^2}\right)^{\varepsilon/2} + (m_L^2)^2 \left(\frac{\mu^2}{m_L^2}\right)^{\varepsilon/2} + (D-1)(m_\gamma^2)^2 \left(\frac{\mu^2}{m_\gamma^2}\right)^{\varepsilon/2} \right]. \quad (12)$$

The constants $Z_j^{(1)}$ are chosen to remove the ε -poles at $D=4$ in the Euler $\Gamma(1-D/2)$ function. They are

$$Z_{m^2}^{(1)} = g \frac{(N+2)}{3\varepsilon}, \quad (13)$$

$$gZ_g^{(1)} = \frac{g^2(N+8) + 108e^4}{3\varepsilon}, \quad (14)$$

$$Z_v^{(1)} = g \frac{N}{2\varepsilon}. \quad (15)$$

From this we obtain in the Landau gauge

$$V^{(0)} = \frac{1}{2}m^2\Phi^2 + \frac{g}{4!}\mu^\varepsilon\Phi^4 + \frac{m^4}{g\mu^\varepsilon}, \quad (16)$$

$$V^{(1,0)} = \frac{\mu^{-\varepsilon}}{8} \left\{ (N-1)m_T^4 \left[-3 + 2\overline{\ln}(m_T^2) \right] + m_L^4 \left[-3 + 2\overline{\ln}(m_L^2) \right] \right. \\ \left. + m_\gamma^4 \left[-5 + 6\overline{\ln}(m_\gamma^2) \right] \right\}, \quad (17)$$

$$V^{(1,\varepsilon)} = -\frac{\mu^{-\varepsilon}}{96} \left((N-1)m_T^4 \left\{ 21 - 18\overline{\ln}(m_T^2) + 6 \left[\overline{\ln}(m_T^2) \right]^2 + \pi^2 \right\} \right. \\ \left. + m_L^4 \left\{ 21 - 18\overline{\ln}(m_L^2) + 6 \left[\overline{\ln}(m_L^2) \right]^2 + \pi^2 \right\} \right. \\ \left. + 3m_\gamma^4 \left\{ 9 - 10\overline{\ln}(m_\gamma^2) + 6 \left[\overline{\ln}(m_\gamma^2) \right]^2 + \pi^2 \right\} \right). \quad (18)$$

A factor $(4\pi)^2$ is absorbed in the definition of \hbar in (10). The function $\overline{\ln}(m^2)$ is defined by

$$\overline{\ln}(m^2) \equiv \ln\left(\frac{m^2}{\mu^2}\right) + \gamma - \ln(4\pi). \quad (19)$$

4. Two-loop contribution

The cancellation of the poles at the two-loop order gives renormalization coefficients $Z_{m^2}^{(2)}, Z_v^{(2)}$ which depend on $\overline{\ln}(m_L), \overline{\ln}(m_T)$ and $Z_\phi^{(1)}$ and a renormalization coefficient $Z_g^{(2)}$ which depends on $\overline{\ln}(m_L), \overline{\ln}(m_T), \overline{\ln}(m_\gamma)$ and $Z_\phi^{(1)}, Z_A^{(1)}$. The results are

$$Z_{m^2}^{(2)} = \frac{(N+2)}{9\varepsilon^2} \left[g^2(N+5) - 18ge^2 + 54e^4 \right] - \frac{1}{6\varepsilon} \left[g^2(N+2) - 8ge^2(N+2) - 6e^4(5N+1) \right], \quad (20)$$

$$gZ_g^{(2)} = \frac{1}{9\varepsilon^2} \left[g^3(N+8)^2 - 18g^2e^2(N+8) + 108ge^4(N+8) + 108e^6(N+18) \right] - \frac{1}{9\varepsilon} \left[g^3(5N+22) - 12g^2e^2(N+5) - 18ge^4(5N+13) + 18e^6(7N+90) \right], \quad (21)$$

$$Z_v^{(2)} = g \frac{N}{6\varepsilon^2} \left[-18e^2 + g(N+2) \right] + ge^2 N \frac{2}{\varepsilon}, \quad (22)$$

$$Z_\phi^{(1)} = e^2 \frac{6}{\varepsilon}, \quad (23)$$

$$Z_A^{(1)} = -e^2 \frac{N}{3\varepsilon}, \quad (24)$$

where a factor $1/(4\pi)^2$ is again absorbed in the definition of \hbar . Then the renormalized two-loop potential in (10) is

$$V^{(2)} = \mu^{-\varepsilon} \sum_{i=0}^2 \sum_{j=0}^2 \sum_{k=0}^2 V_{i,j,k}^{(2)} \left[\overline{\ln}(m_T^2) \right]^i \left[\overline{\ln}(m_L^2) \right]^j \left[\overline{\ln}(m_\gamma^2) \right]^k, \quad (25)$$

where

$$\begin{aligned} V_{0,0,0}^{(2)} = & \frac{1}{432gm_\gamma^4} \left[18m_\gamma^2 \left(g^2m_\gamma^2 [(N-1)(N+5)m_T^4 + 18m_L^2m_T^2 - (2N+13)m_L^4] \right. \right. \\ & + ge^2 \{ 42(m_L^2 - m_T^2)^2(m_L^2 + m_T^2) - 6m_\gamma^2 [(19N-31)m_T^4 + 24m_T^2m_L^2 + 7m_L^4] \\ & \left. \left. - 18m_\gamma^4 [(N-1)m_T^2 + m_L^2] + (19N-36)m_\gamma^6 \} - 108e^4m_\gamma^2(m_L^2 - m_T^2) [(N-1)m_T^2 + 3m_L^2] \right) \right. \\ & + 6g\Phi^2 \{ 7g^2m_\gamma^4 [2(N-1)m_T^2 + (N+8)m_L^2] + 18e^4(7m_L^6 - 7m_L^4m_\gamma^2 + 11m_L^2m_\gamma^4 + 54m_\gamma^6) \} \\ & + \pi^2 \left(g^2m_\gamma^4 [2(N-1)m_T^2 + (N+8)m_L^2] [3(m_T^2 - m_L^2) + g\Phi^2] \right. \\ & + 18ge^2m_\gamma^2 \{ (m_L^2 - m_T^2)^2(m_L^2 + m_T^2) + 3m_\gamma^4 [(N-1)m_T^2 + m_L^2] - 9m_\gamma^6 \} \\ & \left. \left. + 18e^4 \{ -9m_\gamma^4(m_L^2 - m_T^2) [(N-1)m_T^2 + 3m_L^2] + g\Phi^2(2m_L^6 - m_L^4m_\gamma^2 + 5m_L^2m_\gamma^4 + 10m_\gamma^6) \} \right) \right. \\ & + 216(4\pi)^4 ge^2 \left[m_\gamma^2(m_L^2 - m_T^2)^2 I^{(0)}(m_L, m_T, 0) + e^2\Phi^2(m_\gamma^2 - m_L^2)^2 I^{(0)}(m_\gamma, m_L, 0) \right] \\ & + 54ge^2 \left(m_\gamma^2 \{ (N-2)(m_\gamma^2 - 4m_T^2)m_\gamma^2 \xi(m_\gamma, m_T, m_T) \right. \\ & \left. + 2 [m_T^4 - 2m_T^2(m_\gamma^2 + m_L^2) + (m_\gamma^2 - m_L^2)^2] \xi(m_\gamma, m_T, m_L) \} \right) \end{aligned}$$

$$\begin{aligned}
& + e^2 \Phi^2 (m_L^4 - 4m_L^2 m_\gamma^2 + 12m_\gamma^4) \xi(m_\gamma, m_\gamma, m_L) \\
& + 6g^3 \Phi^2 m_\gamma^4 [(N-1)\xi(m_T, m_T, m_L) + 3\xi(m_L, m_L, m_L)] \Big], \tag{26}
\end{aligned}$$

$$\begin{aligned}
V_{1,0,0}^{(2)} = & -\frac{m_T^2}{12gm_\gamma^2} \left(g^2(N-1)m_\gamma^2[-m_L^2 + (N+3)m_T^2 + 2g\Phi^2] \right. \\
& \left. + 6ge^2 \{ 3(m_L^2 - m_T^2)^2 - m_\gamma^2[2m_L^2 + m_T^2(7N-9)] + 5(N-1)m_\gamma^4 \} - 54e^4(N-1)m_\gamma^2(m_L^2 - m_T^2) \right), \tag{27}
\end{aligned}$$

$$\begin{aligned}
V_{2,0,0}^{(2)} = & \frac{1}{72gm_\gamma^2} \left(g^2(N-1)m_\gamma^2 \{ -m_L^2(6m_T^2 + g\Phi^2) + m_T^2 [3m_T^2(N+3) + 4g\Phi^2] \} \right. \\
& - 9ge^2 [-2(m_L^2 - m_T^2)^2 m_T^2 + m_\gamma^2(N-2)(m_\gamma^4 + 6m_T^4) - 6(2N-3)m_\gamma^4 m_T^2] \\
& \left. - 162e^4(N-1)m_\gamma^2(m_L^2 - m_T^2)m_T^2 \right), \tag{28}
\end{aligned}$$

$$\begin{aligned}
V_{0,1,0}^{(2)} = & -\frac{m_L^2}{12gm_\gamma^4} \left(g^2 m_\gamma^4 [-(N+5)m_L^2 + (2N+7)m_T^2 + (N+8)g\Phi^2] \right. \\
& + 6ge^2 m_\gamma^2 \{ 3(m_L^2 - m_T^2)^2 - m_\gamma^2[5m_L^2 + 2m_T^2] + 5m_\gamma^4 \} \\
& \left. - 6e^4 \{ m_\gamma^4 [27(m_L^2 - m_T^2) - 16g\Phi^2] + 3g\Phi^2 m_L^2 (2m_\gamma^2 - m_L^2) \} \right), \tag{29}
\end{aligned}$$

$$\begin{aligned}
V_{0,2,0}^{(2)} = & \frac{m_L^2}{72gm_\gamma^4} \left(g^2 m_\gamma^4 [-3(N+5)m_L^2 + 3(N+8)m_T^2 + (N+17)g\Phi^2] \right. \\
& + 18ge^2 m_\gamma^2 [(m_L^2 - m_T^2)^2 + 3m_\gamma^4] \\
& \left. - 9e^4 \{ 2m_\gamma^4 [27(m_L^2 - m_T^2) - 7g\Phi^2] + g\Phi^2 m_L^2 (4m_\gamma^2 - 3m_L^2) \} \right), \tag{30}
\end{aligned}$$

$$\begin{aligned}
V_{0,0,1}^{(2)} = & \frac{e^2}{6m_\gamma^2} \left\{ -9e^2 m_L^4 \Phi^2 - 3m_\gamma^2 [m_L^4 + m_T^4 - 2m_L^2(m_T^2 + 4e^2\Phi^2)] \right. \\
& \left. + 12m_\gamma^4 [m_L^2 + (N-1)m_T^2 - 6e^2\Phi^2] - m_\gamma^6(4N-9) \right\}, \tag{31}
\end{aligned}$$

$$V_{0,0,2}^{(2)} = \frac{e^2}{8m_\gamma^4} [-18m_\gamma^8 + e^2\Phi^2 (-m_L^6 + 8m_L^4 m_\gamma^2 - 22m_L^2 m_\gamma^4 + 40m_\gamma^6)], \tag{32}$$

$$\begin{aligned}
V_{1,1,0}^{(2)} = & \frac{1}{36m_\gamma^2} \left\{ g(N-1)m_L^2 m_\gamma^2 (3m_T^2 + g\Phi^2) \right. \\
& \left. - 9e^2 [-(m_L^2 - m_T^2)^2(m_L^2 + m_T^2) + 3m_\gamma^2(m_L^4 + m_T^4) - 3m_\gamma^4(m_L^2 + m_T^2) + m_\gamma^6] \right\}, \tag{33}
\end{aligned}$$

$$V_{1,0,1}^{(2)} = \frac{e^2}{4m_\gamma^2} [-(m_L^2 - m_T^2)^3 + 3m_\gamma^2(m_L^4 - m_T^4) - 3m_\gamma^4(m_L^2 - m_T^2) + (N-1)m_\gamma^6], \tag{34}$$

$$\begin{aligned}
V_{0,1,1}^{(2)} = & \frac{e^2}{4m_\gamma^4} \left\{ e^2 m_L^6 \Phi^2 + m_\gamma^2 [m_L^6 - m_L^4(3m_T^2 + 4e^2\Phi^2) + 3m_L^2 m_T^4 - m_T^6] \right. \\
& \left. + m_\gamma^4 [m_L^2(-3m_L^2 + 14e^2\Phi^2) + 3m_T^4] + 3m_\gamma^6(m_L^2 - m_T^2) + m_\gamma^8 \right\}. \tag{35}
\end{aligned}$$

The function $I^{(0)}(m_1, m_2, 0)$ in (26) denotes a non-diverging piece, without the coefficient $(\mu^\varepsilon)^2$, of

$$(\mu^\varepsilon)^2 (4\pi)^4 I(m_1, m_2, 0) = -\frac{2}{\varepsilon^2} (m_1^2 + m_2^2) - \frac{2}{\varepsilon} \left[\frac{3}{2} (m_1^2 + m_2^2) - L_1(m_1, m_2, 0) \right]$$

$$\begin{aligned}
 & -\frac{1}{2} \left\{ L_2(m_1, m_2, 0) - 6L_1(m_1, m_2, 0) + 2m_1^2 \overline{\ln}(m_1^2) \overline{\ln}(m_2^2) + [\overline{\ln}(m_1^2 - m_2^2)]^2 (m_1^2 - m_2^2) \right. \\
 & - 2\overline{\ln}(m_1^2 - m_2^2) \overline{\ln}(m_2^2) (m_1^2 - m_2^2) + 2(m_1^2 - m_2^2) \text{Li}_2\left(\frac{m_2^2}{m_2^2 - m_1^2}\right) + (m_1^2 + m_2^2) [7 + \zeta(2)] \\
 & \left. + \frac{\pi^2}{3} (m_1^2 - m_2^2) \right\}, \tag{36}
 \end{aligned}$$

where

$$L_1(m_1, m_2, m_3) = m_1^2 \overline{\ln}(m_1^2) + m_2^2 \overline{\ln}(m_2^2) + m_3^2 \overline{\ln}(m_3^2), \tag{37}$$

$$L_2(m_1, m_2, m_3) = m_1^2 [\overline{\ln}(m_1^2)]^2 + m_2^2 [\overline{\ln}(m_2^2)]^2 + m_3^2 [\overline{\ln}(m_3^2)]^2, \tag{38}$$

The function $\xi(m_1, m_2, m_3)$ in (26) is defined by

$$\begin{aligned}
 \xi(m_1, m_2, m_3) & \equiv 4 \left(2m_1^2 m_2^2 + 2m_1^2 m_3^2 + 2m_2^2 m_3^2 - m_1^4 - m_2^4 - m_3^4 \right)^{1/2} \\
 & \times \left[L(\theta_1) + L(\theta_2) + L(\theta_3) - \frac{\pi}{2} \ln(2) \right]. \tag{39}
 \end{aligned}$$

where $L(t)$ is the Lobachevsky function, defined by

$$L(t) = - \int_0^t dx \ln \cos x, \tag{40}$$

and the angles are given by

$$\theta_j = \arctan \left[\frac{(m_1^2 + m_2^2 + m_3^2) - 2m_j^2}{(2m_1^2 m_2^2 + 2m_1^2 m_3^2 + 2m_2^2 m_3^2 - m_1^4 - m_2^4 - m_3^4)^{1/2}} \right]. \tag{41}$$

5. Conclusion

As should have become clear from this short review, the superconductive phase transition still poses many interesting problems. In particular, the inclusion of vortex loop fluctuations into the ordinary ϵ -expansion seems necessary to stabilize the renormalization flow in the type-II regime. I hope that after his 60th birthday, Reinhard Folk will have many more active years to contribute to our understanding of this theoretically and experimentally very important phase transition.

Acknowledgements

I am grateful to Dr. F.S.Nogueira for many interesting discussions and to the European Network COSLAB for partial support.

References

1. Folk R., Holovatch Yu., J. Phys. A, **29**, 1996, 3409. 2
2. Folk R., Holovatch Yu., Critical Fluctuations in Normal-to-Superconducting Transition. Correlations, Coherence, and Order, ed. by Shopova D.V., Uzunov D.I. Kluwer Academic/Plenum Publishers, N.Y.-London, 1999, 83–116; cond-mat/9807421. 3
4
5
3. Halperin B.I., Lubensky T.C., Ma S.-K., Phys. Rev. Lett., 1974, **32**, 292; 6
Chen J.-H., Lubensky T.C., Nelson D.R., Phys. Rev. B, 1978, **17**, 4274. 7
4. Lawrie I.A., Nucl. Phys. B, 1982, **200**, 1. 8
5. Coleman S., Weinberg E., Phys. Rev. D, 1973, **7**, 1888; 9
Kleinert H. Limitations on Coleman-Weinberg mechanism. Phys. Lett. B, 1983, **128**, 10
69, (<http://www.physik.fu-berlin.de/~kleinert/106>). 11
6. Ginzburg V.L., Fiz. Tverd. Tela, 1960, **2**, 2031; 12
Sov. Phys. Solid State, 1961, **2**, p. 1824; 13
see also the detailed discussion in Chapter 13 of the textbook L.D. Landau and 14
E.M. Lifshitz, Statistical Physics, 3rd edition. Pergamon Press, London, 1968. 15
Note, however, that the Ginzburg criterion is not applicable to systems with continu- 16
ous symmetry. It specifies where fluctuations drive an order parameter across a barrier 17
separating two phases. In systems with continuous symmetry, the order goes around 18
a barrier. This is much easier, in particular, if the symmetry is high [factor N^2 for 19
 $O(N)$ -symmetry]. The proper criterion is derived in 20
Kleinert H. Criterion for dominance of directional versus size fluctuations of order field 21
in restoring spontaneously broken continuous symmetries. Phys. Rev. Lett., 2000, **84**, 22
286; (cond-mat/9908239) 23
7. Kleinert H. Gauge Fields in Condensed Matter, vol. 1, p. 735–742. Superflow and Vortex Lines. World Scientific Publishing Co., Singapore, 1989, 24
(<http://www.physik.fu-berlin.de/~kleinert/b1>). 25
26
8. Kleinert H., Schulte-Frohlinde V. Critical Properties of Φ^4 -Theories, p. 1–487. World 27
Scientific, Singapore, 2001, (<http://www.physik.fu-berlin.de/~kleinert/b8>). 28
9. Dasgupta C., Halperin B.I., Phys. Rev. Lett., 1981, **47**, p. 1556. 29
10. Als-Nielsen J., Litster J.D., Birgeneau R.J., Kaplan M., Safinya C.R., Lindegaard- 30
Andersen A., Mathiesen S., Phys. Rev. B, 1980, **22**, p. 312. 31
11. de Gennes P.D., Phys. Lett. A, 1969, **30**, 454; 32
Mol. Cryst. Liq. Cryst., 1971, **12**, 193; 33
The Physics of Liquid Crystals, p. 85. Oxford University Press, Oxford, 1993; 34
see also the dual theory in this system: 35
Kleinert H., J. Phys. (Paris), 1983, **44**, 353, 36
(<http://www.physik.fu-berlin.de/~kleinert/102>) 37
12. Schneider T., Khasanov R., Conder K., Pomjakushina E., Bruetsch R., Keller H. 38
Evidence for charged critical fluctuations in underdoped Y-123, (cond-mat/0406691); 39
Schneider T., Khasanov R., Keller H. Evidence for charged critical behavior in the 40
pyrochlore superconductor RbOs₂O₆, (cond-mat/0409398). 41
13. A theoretical alternative would be the uncharged critical point in which $\nu_\lambda = \nu_\xi/2 \approx$ 42
 $1/3$. See [23,24]. 43
14. Kleinert H., Lett. Nuovo Cimento, 1982, **35**, 405, (<http://www.physik.fu-berlin.de/~kleinert/97>). 44
45
15. Bartholomew J., Phys. Rev. B, 1983, **28**, 5378; 46

- 1 Muneshisa Y., Phys. Lett. B, 1985, **155**, 159.
- 2 16. Mo S., Hove J., Sudbo A., The order of the metal to superconductor transition, Phys.
3 Rev. B, 2002, **65**, 104501, (cond-mat/0109260).
- 4 17. Kleinert H., Van den Bossche B., Two-Loop Effective Potential of O(N)-Symmetric
5 Scalar QED in 4-epsilon Dimensions, Nucl. Phys. B, 2002, **632**, 51, (cond-
6 mat/0104102)
- 7 18. Ford C., Jack I., Jones D.R.T., Nucl. Phys. B, 1992, **387**, 373.
- 8 19. Davydychev A.I., Tausk J.B., Nucl. Phys. B, 1993, **397**, 123.
- 9 20. Davydychev A.I., Phys. Rev. D, 2000, **61**, 087701.
- 10 21. Kolnberger S., Folk R., Phys. Rev. B, 1992, **41**, 4083.
- 11 22. Tessman J.-P. Two-loop Renormierung der skalaren Elektrodynamik. Diplomarbeit,
12 Freie Universität, Berlin, 1984, written under my. A pdf file of the thesis is available
13 in the internet: <http://www.physik.fu-berlin.de/kleinert/MS-Tessmann.pdf>.
- 14 23. Kiometzis M., Kleinert H., Schakel A.M.J., Phys. Rev., 1994, **73**, 1975;
15 Herbut I.F., Tesanović Z., Phys. Rev. Lett., 1996, **76**, 4588;
16 de Calan C., Malbuisson A.P.C., Nogueira F.S., Svaiter N.F., Phys. Rev. B, 1999, **59**,
17 554;
18 de Calan C., Nogueira F.S., Phys. Rev. B, 1999, **60**, 4255;
19 Bergerhoff B., Freire F., Litim D.F., Lola S., Wetterich C., Phys. Rev. B, 1996, **53**,
20 5734.
- 21 24. Kleinert H., Nogueira F.S. Charged fixed point in the Ginzburg-Landau superconduc-
22 tor and the role of the Ginzburg parameter κ , Nucl. Phys. B, 2003, **651**, 361 (this
23 paper shows that a charged fixed point is found in $D = 3$ dimensions if $\kappa > 1/\sqrt{2}$).
- 24 25. Only the knowledge of very high orders gives information on such nonperturbative
25 contributions by appropriate resummation. See
26 Hamprecht B., Kleinert H., Variational Perturbation Theory for Summing Divergent
27 Non-Borel-Summable Tunneling Amplitudes, J. Phys. A: Math. Gen., 2004, **37**, 8561,
28 (hep-th/0303163);
29 Phys. Lett. B, 2003, **564**, 111, (hep-th/0302124).
- 30 26. Kleinert H., Phys. Rev. D, 1999, **60**, 085001.
- 31 27. Kleinert H., Path Integrals in Quantum Mechanics, Statistics and Polymer Physics,
32 (variational perturbation theory is developed in chapters 5 and 17). World Scientific.
- 33 28. Kleinert H., Phys. Rev. D, 1998, **57**, 2264;
34 Addendum, 1998, **58**, 107702.
- 35 29. Jasch F., Kleinert H., J. Math. Phys., 2001, **42**, 52.
- 36 30. Kleinert H., Phys. Lett. B, 1998, **434**, 74;
37 Phys. Lett. B, 1999, **463**, 69;
38 Kleinert H., Schulte-Frohlinde V., J. Phys. A, 2001, **34**, 1037.
- 39 31. Lipa J.A., Nissen J.A., Stricker D.A., Swanson D.R., Chui T.C.P., Phys. Rev. B, 2003,
40 **68**, 174518, (cond-mat/0310163).
- 41 32. Kleinert H., Van den Bossche B., Phys. Rev. E, 2001, **63**, 056113.
- 42 33. Strösser M., Larin L.A., Dohm V., Nucl. Phys. B, 1999, **540**, 654.
- 43 34. Brézin E., Wallace D.J., Wilson K.G., Phys. Rev. Lett., 1972, **29**, 591;
44 Phys. Rev. B, 1973, **7**, 232.
- 45 35. Brézin E., Le Guillou J.C., Zinn-Justin J. Phase Transition and Critical Phenomana,
46 edited by C.Domb and M.S.Green, vol. 6., p. 125. Academic Press, New-York, 1976.
- 47 36. For the Ising model, corresponding to $N = 1$, there exists an ε -expansion of the

- equation of state up to three-loop order in 1
 Wallace D.J., Zia R.K.P., Phys. Lett. A, 1973, **46**, 261; 2
 J. Phys. C, 1974, **7**, 3480. 3
37. Kastening B., Kleinert H., Van den Bossche B. Three-Loop Ground-State Energy of 4
 $O(N)$ -Symmetric Ginzburg-Landau Theory above T_c in $4 - \epsilon$ Dimensions with Minimal 5
 Subtraction, Phys. Rev. B, 2002, **65**, 174512, (cond-mat/0109372). 6
38. de Witt B., Phys. Rev., 1967, **162**, 1195. 7
39. Nogueira F.S., Europhys. Lett., 1999, **45**, 612; 8
 de Calan C., Nogueira F.S., Phys. Rev. B, 1999, **60**, 11929. 9
40. Kleinert H., Schakel A., in preparation. 10

Порядок фазового переходу в надпровідний стан

Г.Кляйнарт

Інститут теоретичної фізики, 11
 Вільний університет Берліну,
 Арнімале 146 D–14195 Берлін, Німеччина

Отримано 26 грудня 2004 р.

З нагоди 60-річчя Райнгарда Фолька я привожу короткий опис 12
 теоретичного прогресу в розумінні критичних властивостей над- 13
 провідників. Я відзначаю теоретичні труднощі у виявленні фазового 14
 переходу другого роду в $O(N)$ симетричній моделі Гінзбурга-Ландау 15
 в вимірності $4 - \epsilon$, а також успіх в передбаченні його існування 16
 і визначенні трикритичної точки за допомогою дуальної невтор- 17
 ядкованої теорії. 18

Ключові слова: 19

PACS: 20